


Physical exploration methods and site characterisation



Presenter

Dr Mark Bunch

Cooperative Research Centre
for Greenhouse Gas
Technologies (CO2CRC)

CAGS Summer School
Wuhan, Hubei province, China
1st November 2010

© CO2CRC
All rights reserved


cags

China Australia Geological Storage of CO₂
中澳二氧化碳地质封存

CO₂CRC



Techno-Economic Resource-Reserve Pyramid for CO₂ Storage Capacity



CO2CRC, 2008
Modified from Bachu
et al., CSLF, 2005

Contingent Capacity

- Subset of prospective capacity obtained by considering technical, legal and regulatory, infrastructure and general economic barriers.
 - Value prone to changes as technology, policy, regulations and/or economics change.
- Corresponds to “Reserves” as used in energy and mining industries



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存




© CO2CRC
All rights reserved



Criteria for Site Selection (In All Geological Storage Systems):

- Capacity (what volume of CO₂ can the rock hold?)
- Containment (can we keep the CO₂ in the rock?)
- Injectivity (can we put the CO₂ into the rock?)
- Other (Economic, Regulatory, Risk, Legal, Community...)





Geological storage of CO₂

What do we need?

SEAL ROCK – variable porosity, low permeability, e.g. claystone

RESERVOIR ROCK – porous, permeable, e.g. sandstone



How do we know what's down there? (Sources of geological data)

- Geophysics
 - Seismic
- Drilled wells
 - Rock (core; drill cuttings)
 - Wireline logs
- Analogs
 - Outcrops
 - Modern depositional environments
- Integrated models



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved




Onshore seismic




What is seismic?



Photo courtesy Philips Research



Animation courtesy US Department of Energy
Collecting seismic data on land



Ultrasound examination during pregnancy

Seismic imaging uses reflected sound waves to create a picture of underground rock formations.



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存




© CO2CRC
All rights reserved



Seismic section

SW

NE



cags


China Australia Geological Storage of CO₂
中澳二氧化碳地质封存

CO₂ CRC

© CO₂CRC
All rights reserved



Geological model




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved





3D layered Earth model



Well data


Core is very important



cags


China Australia G
CO₂ CRC
中澳二氧

Wireline well logging




- Gamma ray, neutron, density, resistivity and caliper log
- NMR
- ECS (elemental capture spectroscopy)
- FMI (image log)
- Sonic Scanner
- Formation tester
- 3D VSP

Wireline well log data



← 1 – 10's of km. →



Wireline log data
(correlation between wells)




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



Studying the rocks



Identifying the likeliest depositional system

Transgressive Shoreface Model:~ from Buffin (1989)

Regressive Braided Fluvial (Faulkner 2001)





Figure 17. Facies distribution and depositional environments throughout the Port Campbell Embayment: during the Upper Waarre Formation.



Possible but unlikely

- Flaxmans A shale is now interpreted to never be time equivalent of Waarre C.

Most likely model imposes modeling constraints:

- channel orientation is grossly N-S.
- but ranges from SE-NW.




Integrated data sources




Core data


- Dep. Env.
- Poro/Perm
- Stratigraph



Outcrop analogs




Seismic



Static model


Qualitative basin ranking



Gippsland Basin, Southeast Australia

The Problem:

- New brown coal developments in Latrobe Valley, Victoria
- CO₂ emissions up to 50 Million tonnes/year




Potential Solution:

- Offshore Gippsland Basin
- Existing oil and gas fields (once depleted)
- Deeper saline formations

C. Gibson-Poole

Map of the Kingfish field, Gippsland Basin



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存




© CO2CRC
All rights reserved




Seismic cross-section and geological interpretation: Kingfish field, Gippsland Basin

Seismic Cross-Section (line G92A-3074A) between Kingfish-3 and Roundhead-1




C. Gibson-Poole


Modelling CO₂ migration pathways: offshore Gippsland Basin



Modelling CO₂ migration pathways: Kingfish field, Gippsland Basin






Modelling storage potential: onshore Gippsland Basin




Bunch et al., 2010

Comparison of storage litho-facies



Bunch et al., 2010
(GHGT-10)

Skeleton of the numerical geo-model



cags


China Australia Geological Storage of CO₂
中澳二氧化碳地质封存

CO₂ CRC

© CO₂CRC
All rights reserved



Storage litho-facies populated across model



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存




© CO2CRC
All rights reserved



Permeability

(measure of fluid connectivity)


$$K = \frac{Q \mu L}{A (P_1 - P_2)}$$



Measured in Darcies (D) or millidarcies (mD)




Porosity and permeability



Porosity is the storage space shown by the blue spaces in this thin slice through a reservoir sandstone.

Permeability is the ability of the rock to allow fluid flow from pore to pore. Permeability is strongly affected by the geometry of the pore throats (red circles).

Permeability is main control on injectivity



Injectivity

$I_{v/t}$ depends on $A * P_i * k$

$I_{v/t}$ = Injection rate

A = Area (of wellbore in contact with formation)

P_i = injection pressure (below frac)

k = permeability

($I_{v/t}$ is proportional
to number of wells)



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



Injectivity / Pressure Considerations:

- Injection of fluids (eg CO₂) causes reservoir pressure build up
- In depleted fields, pressure build-up may be neutral or beneficial
- In both depleted fields and saline aquifers, pressure must remain below fracture pressure
- In low permeability reservoirs this may limit economic storage capacity due to decreased injection rate, requiring more wells
- Injection in saline formations may displace saline fluids & increase risk of possible mixing with a freshwater system
- Drilling pressure relief (water production) wells is a possible solution




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存




© CO₂CRC
All rights reserved




Depleted Field (Pressure v. Time)



Saline Aquifer (Pressure v. Time)




Modelling CO₂ storage capacity – the effect of injectivity on economics




Allinson & Paterson, 2009

Containment

Buoyancy vs. capillary pressure



$$P_b > P_c$$




$$S_w = 100\%$$

$$P_b < P_c$$




Mercury porosimeter




$$P_m > P_c$$




Mercury Capillary Pressure Experiment



Mercury Capillary Pressure Experiment



Mercury Capillary Pressure Experiment




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存




© CO2CRC
All rights reserved



Mercury Capillary Pressure Experiment



Mercury injection capillary pressure



Mercury (Nonwetting-phase) Saturation



China Australia Geological Storage of CO₂

中澳二氧化碳地质封存






© CO2CRC
All rights reserved



Site characterisation process

- Build detailed reservoir model using current state of the art modelling packages
- History match with actual production data to validate model.
- Predict future trend.




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



Monitoring the injected CO₂



Measuring the temperature and pressure, recording sound waves and detecting chemical and electrical changes



Seismic monitoring



cags

China Australia Geological Storage of CO₂
中澳二氧化碳地质封存


Image, CO2CRC

CO₂ CRC

© CO2CRC
All rights reserved




The Sleipner 4D seismics until 2006




Vertical seismic profile monitoring

Zero offset VSP




The source is close to the well.

Offset VSP




The source is some distance from the well.

Deviated Well VSP (ZVSP)



Multiple geophones are located in the well, with multiple source positions.

Walkaway VSP (WVSP)




The source is moved away from the well and signals are generated at regular spacings.


- VSP tomography
- High Resolution Travel time
- Microseismic surveys (measures creaks in the subsurface)



Downhole geochemical monitoring



Model predictions




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



Cross-well electromagnetic tomography



Sibblmønger
S&TR, May 2005

Crosswell electromagnetic (EM) imaging of CO₂ sequestration at the Lost Hills field in central California—an enhanced oil recovery site operated by Chevron USA




China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved




Model validation through new data



Pre-production GWC,
2020 m S.S.

Current GWC, 1990 m S.S.
(2040 m R.T.)

1. Gas cap
2. Trapped gas ($S_{gr}=0.25$)
3. Aquifer (partial)



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



Other monitoring options...

- Microseismic activity, deep and shallow
- Pressure sensors in monitoring well
- InSAR to measure mm-to-cm scale surface deformation



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



Acknowledgement

The authors would like to acknowledge the funding provided by the Australian Commonwealth through the CRC Program, and by both industry and state government partners to support CO2CRC research.



CRC FOR GREENHOUSE
GAS TECHNOLOGIES



Thank you

www.co2crc.com.au



China Australia Geological Storage of CO₂
中澳二氧化碳地质封存



© CO2CRC
All rights reserved



CO2CRC Participants



Supporting Partners: The Global CCS Institute | The University of Queensland | Process Group | Lawrence Berkeley National Laboratory

Established & supported under the Australian Government's Cooperative Research Centres Program

